PRIZE ESSAY

MICROBIOME

Poised for tissue repair

Skin microbes interact with the immune system to aid wound healing

By Oliver Harrison

he immune system acts as a formidable guardian of health, protecting and restoring tissue function during infectious and environmental challenges. To date, our understanding of host immunity stems from models of inflammation and infection with pathogenic microbes. However, the vast majority of microbial-immune encounters occur as a symbiotic relationship with the commensal

microbiota, the trillions of microorganisms that inhabit our barrier tissues, including the gastrointestinal tract and skin. Indeed, far from being ignored by the host, commensal microbes promote the development, education, and

function of the mammalian immune system (1). In return, immune responses to commensal microbes, particularly those mediated by the adaptive immune system, act to bolster epithelial fortifications for antimicrobial function, restraining systemic translocation of both commensal and pathogenic microbes (2).

A striking feature of immune responses to commensal microbes present at barrier surfaces is the uncoupling from inflammation and maintenance of tissue integrity at both the onset and effector phases of the response. This form of homeostatic immunity to commensal microbes raises several questions. In particular, to what extent do the conventional rules of adaptive immunity to pathogens apply to these noninvasive microbes? And, how do commensal-specific T cells sense and respond to environmental stresses during tissue injury?

As a postdoctoral fellow in Yasmine Belkaid's laboratory at the National Institutes of Health, I sought to address these questions, investigating the mechanisms by which commensal-specific T cell responses are mounted and their role in tissue homeostasis. Initial efforts to identify, track, and profile commensal-specific T cells led us to uncover

Center for Fundamental Immunology, Benaroya Research Institute, Seattle, WA, USA Email: oharrison@benaroyaresearch.org

three aspects of host immunity specifically engaged by commensal microbes resident at the skin surface: a nonclassical major histocompatibility complex (MHC) restriction, a hybrid differentiation profile, and a direct contribution to wound repair.

NONCLASSICAL IMMUNITY TO **COMMENSAL MICROBES**

Despite the overwhelming quantity of proteins potentially produced by the microbiota, only a smattering of microbe-derived

> antigens and epitopes recognized by commensal-specific T cells have been identified. hampering mechanistic insight into the biology of these cells (3-6). For this reason, we began by exploring the antigen-specificity of

commensal-specific T cells resident within the skin, using a model of neocolonization with the common human skin commensal Staphylococcus epidermidis (7).

A combination of in vitro screening and in silico epitope prediction revealed that commensal-specific CD8+ T cell responses were coordinated by the noncanonical MHC1b molecule, H2-M3, presenting *N*-formylmethionine (fMet)-containing peptides (8). These findings demonstrated


that fMet peptides, a canonical class of bacterial antigens known to activate the innate immune system, also induced a tailored adaptive immune response to noninvasive bacteria. They also highlighted the distinct contribution of nonclassical MHC molecules to the cross-talk that occurs between microbiota and the immune system. Identification of H2-M3-binding fMet peptides facilitated the generation of peptide:MHC tetramers and T cell receptor (TCR)-transgenic mice, enabling the tracking and profiling of commensal-specific CD8+ T cells within skin tissues during homeostasis and inflammation. Using these tools as complementary strategies to track commensal-specific CD8+ T cells in vivo, we determined that these T cells display a tissue-resident memory profile, surviving locally as long-term sentinels of the skin barrier.

POISED IMMUNITY TO **COMMENSAL MICROBES**

To determine how commensal-specific CD8⁺ T cells were educated by the local skin microenvironment, we assessed their cytokine production and effector potential. Commensal-specific CD8+ T cells polarized toward an unusual hybrid state, highly distinct from those commonly invoked by pathogen infections. Specifically, CD8+ T

Commensal-specific CD8⁺ T cells harbor a poised type-2 transcriptome during homeostasis

Tissue injury licenses poised type-2 immunity, including translation of interleukin-13 (IL-13) protein by commensal-specific CD8⁺ T cells, thereby promoting tissue repair.



NOSTER Science


MICROBIOME

PRIZE

cells coexpressed two normally mutually exclusive transcription factors, retinoic acidrelated orphan receptor- γt (ROR γt) and GATA-binding protein 3 (GATA-3), that are essential for type-17 immunity to fungi and type-2 immunity to parasites, respectively. We next investigated the impact of such coexpression by characterizing the epigenetic and transcriptional landscape of these cells.


Unexpectedly, we identified accessible chromatin and mRNA expression for type-2 immunity cytokines, *Il5* and *Il13*. By contrast, although competent for *Il5* and *Il13* mRNA expression, these cells did not subsequently translate either interleukin-5 (IL-5) or IL-13 proteins. Thus, in healthy skin tissue, commensal-specific T cells adopt a distinct differentiation profile, which we have termed "poised type-2 immunity" (9).

A mechanistic understanding of the posttranscriptional machinery that regulates this immune axis remains a focus of ongoing research. However, given the fundamental role



CO-OPTING LESSONS FROM COMMENSAL-SPECIFIC IMMUNITY TO UNDERSTAND DISEASE

By identifying a tissue checkpoint that is reliant on the plasticity of tissue-resident com-



Confocal imaging of the T cell response to S. epidermidis colonization of the skin. Foxp3, green; DAPI, blue; CD8 α , red.

of the skin as a protective barrier, we hypothesized that poised type-2 immunity might be influenced by environmental insults, including tissue injury. We determined that it could be unleashed through stimulation with local alarmins, factors released during skin inflammation, triggering rapid protein translation of type-2 cytokines by commensal-specific CD8⁺ T cells locally within injured skin tissue. Thus, induction of type-2 cytokine mRNA expression and subsequent protein translation can be temporally decoupled in commensal-specific CD8⁺ T cells, allowing for pleiotropic functions under homeostatic and inflammatory conditions.

COMMENSAL-SPECIFIC T CELL PLASTICITY AIDS TISSUE REPAIR

Given the anatomical proximity of commensal-specific $CD8^+$ T cells to the skin epithelium, and their remarkable functional mensal-specific T cells, we extend our understanding of a fundamental but poorly understood form of immunity. Our findings highlight the exquisite specificity that underlies the interplay between commensal microbes and the immune system as a key element of tissue homeostasis and repair. We, like others, hypothesize that perturbations in this symbiotic relationship are likely triggering events for the onset and perpetuation of chronic inflammatory disorders of barrier tissues. Further understanding the contribution of commensal-

specific immunity to tissue physiology may tailor future therapies toward critical immune-microbe interactions driving chronic inflammation and disease.

REFERENCES AND NOTES

- 1. Y. Belkaid, O. J. Harrison, Immunity 46, 562 (2017).
- 2. K. Honda, D. R. Littman, *Nature* 535, 75 (2016).
- Y. Cong, T. Feng, K. Fuji-hashi, T. R. Schoeb, C. Ó. Elson, Proc. Natl. Acad. Sci. U.S.A. 106, 19256 (2009).
- 4. Y. Yang et al., Nature **510**, 152 (2014).
- M. Xu et al., Nature 554, 373 (2018).
- 6. E. Ansaldo et al., Science **364**, 1179 (2019).
- 7. S. Naik et al., Nature **520**, 104 (2015).
- 8. J.L.Linehan et al., Cell 172, 784 (2018)
- 9. O. J. Harrison et al., Science 363, eaat6280 (2019).

ACKNOWLEDGMENTS

I am extremely grateful to my mentor, Y. Belkaid, with whom I am lucky to have trained and benefitted from her vision, drive, and mentorship for several years. I also thank the members of the Belkaid laboratory, past and present, and our collaborators for their generosity in advice and friendship.

10.1126/science.abc5618



GRAND PRIZE WINNER Oliver Harrison

Oliver Harrison re-

ceived an undergraduate degree from the University of Bath and

a Ph.D from the University of Oxford. After completing his postdoctoral fellowship at the National Institutes of Health, Oliver started his laboratory in the Center for Fundamental Immunology at Benaroya Research Institute in 2019. His research investigates how T and B cell responses to commensal microbes promote barrier tissue integrity and repair and how this goes awry in disease.



FINALIST Mariana X. Byndloss

Mariana X. Byndloss received her D.V.M. and a Ph.D. from Universidade Federal de Minas Gerais in Brazil. After

completing her postdoctoral fellowship at the University of California, Davis, Mariana started her laboratory in the Department of Pathology, Microbiology, and Immunology at Vanderbilt University Medical Center in 2018. Her research aims to understand how inflammationmediated changes in gut epithelial metabolism lead to gut dysbiosis and increased risk of infectious gastroenteritis by *Salmonella typhimurium* and noncommunicable diseases, namely obesity-associated cardiovascular disease and colon cancer. www.sciencemag. org/content/369/6500/153.1



FINALIST Chun-Jun (CJ) Guo

Chun-Jun (CJ) Guo received undergraduate degrees from Fudan University and a Ph.D. from the University of

Southern California. After completing his postdoctoral fellowship at the University of California, San Francisco, and Stanford University in the Fischbach group, CJ started his laboratory in the Jill Roberts Institute for Research in Inflammatory Bowel Disease at Weill Cornell Medicine in 2018. His research uses CRISPR-based microbial genetics, state-of-the-art analytical chemistry, and healthy and diseased mouse models to mechanistically dissect how microbiome-derived genes, pathways, and metabolites affect host biology. www.sciencemag.org/content/ 369/6500/153.2

Downloaded from http://science.sciencemag.org/ on July 10, 2020

VANDERBILT UNIVERSITY MEDICAL CENTER; COURTESY OF CHUN-JUN GUO

PHOTO: (CLOCKWISE FROM LEFT) 0.J. HARRISON ET AL. (9); COURTESY OF OLIVER HARRISON;



Poised for tissue repair

Oliver Harrison

Science **369** (6500), 152-153. DOI: 10.1126/science.abc5618

 ARTICLE TOOLS
 http://science.sciencemag.org/content/369/6500/152

 RELATED CONTENT
 http://science.sciencemag.org/content/sci/369/6500/153.1.full http://science.sciencemag.org/content/sci/369/6500/153.2.full http://stm.sciencemag.org/content/sci/ansmed/11/518/eaau4760.full http://stm.sciencemag.org/content/sci/ransmed/11/507/eaau9356.full http://stm.sciencemag.org/content/sci/ransmed/11/502/eaan5662.full http://stm.sciencemag.org/content/sci/ransmed/11/502/eaan5662.full http://stm.sciencemag.org/content/sci/ransmed/11/502/eaan5662.full http://stm.sciencemag.org/content/sci/ransmed/11/502/eaan5662.full http://stm.sciencemag.org/content/sci/ransmed/11/502/eaan5662.full http://stm.sciencemag.org/content/sci/ransmed/11/502/eaan5662.full http://stm.sciencemag.org/content/sci/ransmed/11/502/eaan5662.full

 REFERENCES
 This article cites 9 articles, 3 of which you can access for free http://science.sciencemag.org/content/369/6500/152#BIBL

 PERMISSIONS
 http://www.sciencemag.org/help/reprints-and-permissions

Use of this article is subject to the Terms of Service

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2020 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works